

priVet Poultry Booster 850 g | 850 g

For upgrading poultry feed at any age

Product number	4500
Weight	1kg
Length	99mm
Width	99mm
Height	222mm



Product description

Complementary feed (powder) for poultry

For upgrading poultry feed at any age

priVet Poultry Booster is designed for healthy growth with an intensive supply of natural minerals from the HuminoMin® complex to the bones. **priVet Poultry Booster** can optimize feed intake by improving the metabolic rate, which can lead to faster growth and corresponding weight gain. At the same time, joint condition and bone structure are supported and the shell strength and laying performance of laying hens can be improved. The meat quality and weight development of animals for fattening can also be improved by a natural supply of minerals and trace elements.

Composition: linseed expeller, brewers' yeast, maerl, sunflower seed expeller, peat, grass, herbs, legume plants, seaweed meal, grape pips meal, brewers' yeast cell walls (MOS), apple pomace, parsnip (ground), aronia pomace, blackcurrant pomace, eggshell powder, carrot pomace, celery tuber (ground), moringa leaves (ground)

Additives/kg: Technological additives: clinoptilolite of sedimentary origin (1g568) 0.6 g. bentonite (1m558i) 0.2 g. Total quantity of clinoptilolite of sedimentary origin from all sources shall not exceed the maximum content of 10000 mg/kg. The total amount of bentonite must not exceed the permitted maximum level in complete feedingstuff of 20000 mg/kg of complete feedingstuff.

Analytical constituents: crude protein 35.2%, crude fat 7.4%, crude fiber 6.8%, crude ash 7.8%, calcium 0.82%, phosphor 0.98%, sodium 0.14%, methionine 0.62%, lysine 1.51%

Feeding recommendation: Depending on growth requirements and feed quality, add up to 20-40% of the total daily ration to the feed. If required, e.g. in stressful situations (moulting and transportation), the feeding quantity can be

doubled. 1 top. corresponds to ca. 3 g. 1 tbsp. corresponds to ca. 9 g.

The simultaneous oral use with macrolides shall be avoided. The simultaneous use with robenidine shall be avoided. The simultaneous use with coccidiostats other than robenidine is contraindicated with level of bentonite above 5000 mg/kg of complete feed.

Product properties

Phase of life	Junior, Erwachsener, Senior
Product type	Ergänzungsfuttermittel (Pulver)
Anwendungsart	Fütterung, innere Anwendung
Animal categories	Geflügel, Huhn, Pute, Wachtel
Composition	Lein, Bierhefe, Algenkalk, Sonnenblume, Moor, Gras-, Kräuter-, Leguminosenpflanzen, Seealge, Traubenkern, Bierhefe, Apfel, Pastinak, Aronia, Schwarze Johannisbeere, Eierschale, Möhre, Sellerie, Moringa, Klinoptilolith
Special Requirements	Knochen, Gelenke, Wachstum, Stoffwechsel, Futteraufnahme, Gewicht, Mineralstoffe, Spurenelemente

More images